The Agricultural College Is a Friend to the Farmer



ITS SOLE AIM IS TO AID AGRICULTURISTS. VIEW OF OREGON AGRICULTURAL COLLEGE AT CORVALLIS, OR.

Now Is Time for Sweet Pea Growers to Hustle

"The sowing of this crop can good garden soil, providing this trouble from over-shading can occur. A row running north and south, right out in full sunlight, is to be preferred. It is true that sweet peas often do well even planted along the north side of a house, but speaking of av-erage conditions, the open is to be

"The soil should be carefully pre-pared, spading to a depth of at least a foot, and turning in, in the lower six inches of this soil, a liberal six inches of this soil, a liberal amount of well-decomposed barn-yard manure. It is best to prepare the soil in this manner during the Fall, and, if possible, to plant the seed at that time. Should planting not prove feasible, and in case it is delayed until Spring, the top six inches of the soil should be stirred over again with a spade before plant-

ing.
"As soon as the soil is in good As soon as the soil is in good working condition, the sweet peas can be set out, or sown. For sowing the seed, dig a small trench or furrow, about 4 or 5 inches deep, smooth the surface of the soil in the bottom of this trench and sow the seed evenly, using about one ounce of seed to 15 or 18 feet of the row. Then cover with two inches of soil. When the seedlings are four or five inches high, one can thin out so that the plants will remain about six inches apart in

the row."

At this time one should have calalogues from the various seedsmen at hand, and he will be able to use these publications to aid him in selecting pathications to aid film in selecting his varieties of sweet peas. Many people prefer to plant sweet peas in mixtures. In this case some of the most satisfactory varieties can be ob-tained, and at that somewhat more cheaply than could the named va-

rieties in separate packets.

There are several places in town where sweet pea seed can be purchased, and the prices run from 5 cents a packet, for a liberal amount of seed, up to much higher prices for the newer varieties, which are offered

o'clock Saturday afternoon, February 6, Carryalls left the college campus at 1:30 so that all who were interested in the demonstrations could be taken to the ground with out charge. The demonstrations were under the immediate charge of

A page of interesting items from the Oregon Agricultural College at Corvallis will alternate in the farm weekly with a page of news notes from the Washington State College at Pollman. This will afford an interchange of views from the two big agricultural colleges of the Northwest that should prove of benefit to the reader, for the institutions deal with similar problems,

Professor H. D. Scudder, of the agronomy department.

Umatilla Sub-Station Issues Report of Work

"I' is time now to consider the planting of sweet pens," says Soll and climatic conditions pre-Professor A. L. Peck, landscape gar- ation project were considered by the dener of Oregon Agricultural Col- reclamation service to be better adapted to the production of fruit be carried on either in the Fall of than to other crops. As a result of the year, or at this time in early this early decision, the land was di-Spring. The peas should be sown in vided into small units with a view to the development of small intensively ground is not subject to heavy soak-ing of water from now on. It should be planted in such situation that no ment station made by Superintendent port of the Umatilia branch experi-ment station, made by Superintendent Ralph W. Allen and issued by the Oregon Agricultural College experi-ment station under the direction of Director A. B. Cordley. The report tells further how and for what pur-pose the branch station was estab-lished, how it is maintained and man-aged, how it views the leading feat-ures of its activities since its estabures of its activities since its estab-lishment. The report says further:

"The predominating soil type on the Umatilla project, upon which the station is located, is sand, ranging in texture from coarse to fine. Approximately half of this area is of coarse sand and the remainder ranges in character from medium to fine. The higher land that lies back from the Columbia River is mostly of finer texture. The soils are markedly de-ficient in organic matter and nitro-The physical character of this land renders the duty of irrigation water very low. From a soil stand-point, the correction of these two difficulties, which are among the prin-cipal factors influencing crop pro-duction, is of the utmost importance.

"Climatic conditions of this district are very congenial for crop growth. They are a rare combination of dry atmosphere and dry weather. The effect is comparatively long growing

seasons and mild, open Winters."

Conditions on this tract are such as render necessary scientific investigation in eradication of alkali, secur-ing stands of crops, crop rotations, and the economical use of water and

Reports of experiments with truck crops, strawberries, cane fruits and tree fruits as carried on in field plats are given in the new bulletin, free copies of which may be had by all interested persons are recovered. terested persons upon request ad-dressed to the Oregon Agricultural

Students Give Book Fund.

students of 1915 have requested the college authorities to accept the special fund raised for this at the Oregon Agricultural College, This action by the solid and stump burning, stump pulling and stump burning, stump pulling and stump burning, stump pulling and stump blasting. Each process was handled by experts who showed six different processes. The demonstration was abort distance from Corvallis at 2 o'cleck Saturday afternoon, Pebruary 6 Carryalls left the approach of the approach of the special fund raised for this college and of the valuable information and training which they secured during their short course work. This action was in line with precedents established by other above that the lamb takes all the milk caked at the course classes at the course classes at the course work. This action was in line with precedents established by other above the course classes at the course work. The demonstration took place on a farm a short distance from Corvallis at 2 o'cleck Saturday afternoon, Pebruary 6 Carryalls left the approach of the second the course work. The demonstration took place on a farm a short distance from Corvallis at 2 o'cleck Saturday afternoon, Pebruary 6 Carryalls left the approach of the second the course work. The demonstration took place on a farm a short distance from Corvallis at 2 o'cleck Saturday afternoon, Pebruary 6 Carryalls left the course of the course work. The demonstration took place on a farm a short distance from Corvallis at 2 o'cleck Saturday afternoon, Pebruary 6 Carryalls left the course work. The course classes at the course work. The course work of the course work of the course work of the course work. The course work of the course

ily for the use of short course students, they are also accessible to the degree and the vocational students and to the college faculties. The list and to the college faculties. will be selected and purchased by Dean A. B. Cordley, of the school of agriculture, and Mrs. Ida Kidder, librarlan.

The class of 1915 elected J. H. ines, of Waldport, president, "be-Glines, of Waldport, president, cause," said the members, "he has attended two short courses before and knows the 'ropes,'". Mrs. J. M. Waldrip, who owns and manages a farm on San Juan Island in Puget Sound, was elected secretary, and H. F. Hawkins, of Salem, was elected treasurer. Mr. Hawkins is now managing the dairy end of a 1600-acre ranch near Dallas. P. C. Burt, of Bend, was chosen to solicit the special funds for the library donation.

Report of John Jacob Astor Branch Station

N 1913 an agricultural experiment N 1913 an agricultural station was established in Clatsop County for the purpose of investigating and demonstrating agricultural possibilities of reclaimed swamp land and logged off land. The biennial report of this station, known as the John Jacob Astor branch experiment station, has been issued by the Agri-cultural College Experiment Station under the direction of Director A. B. Cordley. In this report, issued as an experiment station bulletin, are summed up the principal facts of establishment and maintenance of the station and the result of its operation

the John Jacob Astor branch station is situated four miles from Astoria on Young's River. Of the 70 acres comprising the farm, 50 acres are tide land and 20 acres logged-off hill land. This represents the average types of land in Clatsop County and the methods and results secured in its management and use will prove of much interest to owners of similar

Methods of draining and clearing the tide lands of their dense growth of rushes, reeds and swamp grasses are summarized in the report. Some of these methods were quite different from those generally followed in clearing these lands and some very clearing these lands and some very excellent results have been secured by them. In addition to this the bulletin describes the cropping systems that were employed successfully in connection with the clearing methods.

Care of Spring Lambs.

"The Spring of the year is the shepherd's harvest. After the long Winter of ceaseless toll and care the lambing time is the joy and the fruit of the shepherd's labors. To him it is the greatest pleasure to welcome each new-born babe, whether it be single, twin or triplet. But like the grain harvest, lambing time is not a time of case. The successful shep-

time of ease. The successful shepherd is very skillful at this time and uses his best judgment in feeding and caring for the flock."

So says Professor O. M. Nelson, specialist at the Oregon Agricultural College in sheep production. He believes that a little grain should be given the ewes a month or so before lambing. Small amounts should be used in beginning the feed, which may consist of one and one-half parts of oats to one part bran fed with

production rests upon marketing, yet this phase of the subject has been given but scant attention while methods of production have engaged the serious consideration of Oregon grow-ers for many years," says Professor A. G. Bouquet, head of vegetable gardening at the Agricultural College.

"Every plan and all the activities of growers are based upon their effects on the sale of his products. Since success in marketing each product by the route that brings the mighest net returns to the grower depends upon having the kind and quality of product demanded by accessible markets, the successful market gardener will always take his market demands into consideration in planning his garden and selecting his crops. It is what he markets, not what he grows, that determines the success of the gardener.'

Time Now to Set Hens to Get Winter Layers

T HE latter part of March or first of April is the best time that Eastern Oregon poultry raisers can select for hatching next Winter's layers, according to Professor A. Q. Lunn, of the Oregon Agricultural College poultry department. Eggs for these hatchings would, of course, be set from the first to the middle of March.

"A difficulty in producing pullets that will lay during the Winter months is the problem of preventing them from beginning to lay before cold weather," continued Mr. Lenn, "If they mature and begin laying earlier than this they will likely molt when they should be producing egg.
When cold weather catches the pullets in molt they need all their por-

lets in molt they need all their powers to maintain bodily vigor and usually will stop laying until the warm days of Spring come.

"If pullets molt in Winter it is probably because they are hatched too early in the season. They reach maturity while it is still warm, begin laying and then go into the molt. This is where one of the greatest troubles lies.

"It is a mistake also to think that pullets hatched in Winter will make the best Winter layers.

"A Leghorn hen requires on the average seven months to mature to the point where it will begin to lay. Heavy breeds of chickens require one or two months additional. Leghorn chickens hatched to mature the

Heavy breeds of chickens require one or two months additional. Leghorn chickens hatched soon after the middle of March, if they make normal development, should begin to lay in October when their chances of moting will be reduced to the minimum. "Chicks that are hatched in late Spring or early Summer are not apt to thrive well on account of the scarcity of proper forage and because of

ity of proper forage and because of other influences that tend to keep them from growing into vigorous and healthy fowls and good layers."

Rural Life Supports Urban.

"Out of the agricultural colleges must come the solution of the great and serious problem of building a rural life to stand the strain of car-ing for the congested cities of the future," said Walter M. Pierce, of the Oregon Agricultural College Board of

Regents.
"More than that, it will be the peculiar mission of the agricultural colleges to make farm life so profitable and attractive that rural settle-

Hog Breed Small Factor.

In starting hog raising growers need not be concerned very seriously about the breed of hogs they select, according to the O. A. C. animal husbandry department. Each of the three principal breeds in Oregon— Poland China, Berkshire and Duret Jersey, has its points of superiority and inferiority, and there is said to be about as much difference in [ad]. be about as much difference in individuals as between breeds. Some run more to the fat hog and lard type and others more to the bacon type, while some are rather earlier maturing than others. But the main consideration in selecting the breed is to choose the one that suits the breeder's ideas and purposes best, and then stick to it and build it up by careful selection and proper management.