

FIRE BLIGHT OF APPLE AND PEAR

Text of Bulletin by H. S. Jackson of the Oregon Agricultural College and Experiment Station.

Fire Blight is the most serious of all the diseases which attack the pear and apple. It is a contagious disease of bacterial origin which, under proper conditions, may attack any part of the tree, besides the pear and apple, the quince, wild crab-apple, hawthorn, mountain ash, serviceberry and some other pomaceous trees are subject to attacks of this disease.

Cause of the Disease.
Fire Blight is caused by a minute organism belonging to the group of bacteria and known technically as *Bacillus amylovorus*. It is a rod-shaped, motile germ which divides very rapidly by simple fission and is found in immense numbers in the diseased tissue. The germs are so minute that they measure only about 1-25,000 of an inch in length and are visible only under the highest magnification of the compound microscope.

Proofs of the Bacterial Nature of the Disease.
Myriads of germs are present in all freshly blighted portions of the tree and in the sticky ooze from exuding cankers. Pure cultures of the organism may be obtained from diseased tissue and cultivated for generation after generation on suitable culture media. Healthy tissue inoculated with germs from a pure culture will become characteristically diseased. The germs are found in abundance in the tissues so inoculated and when grown in pure culture and compared with germs from naturally infected tissue, and with those used to produce the artificial infection, are found to be identical.

Part of the Tree Affected.
The germ lives almost entirely in the sappy portion of the bark, though in some vigorous-growing varieties of pears the germs have been known to invade the sap wood to a limited extent. When inoculated into the growing tips of branches, into the blossom or the fruit, the germs are found generally throughout the tissues.

Distribution.
Fire Blight occurs in more or less severity in nearly all parts of the United States where pears and apples are grown. It was first reported in the Hudson River valley, N. Y., about 1792 and was doubtless indigenous in the eastern United States. The disease has gradually spread westward as the country became settled and according to Waite became known in California about 1895-1897. According to the same author it has been known in British Columbia for almost the same period. It is probable that the disease has progressed westward along two rather distinct routes—one north into Idaho, Washington and British Columbia, the other south into California.

In Oregon Fire Blight has appeared in two general localities—one in the southwestern part of the state, including the Rogue River valley, the other in the northeastern part. The former region has doubtless become infested by the spread of the disease north from California, while the eastern part of the state has been infected through Idaho or Washington along the northern route.

Life Cycle of the Blight Organism.
Beginning in the spring the first apparent damage produced by the disease in an infected orchard is the blighting of the blossoms. Infection is brought about by insects, principally bees, which have visited a case of hold-over blight and become covered with the organisms contained in the sticky exudation, inoculating the flowers in their search for nectar. The organisms divide and multiply in the nectar and are able to enter the living tissues through the unprotected nectaries. Having entered the tissues they quickly blight the blossoms, pass down the blossom stem and into the fruit spur, killing the tissues and cutting off the leaves from water supply causing them to shrivel and dry, thus producing "fruit spur blight". The latter occurs several weeks after blossom infection. In very serious

cases nearly all the fruit spurs may be blighted in this way and the trees set no fruit. Usually the germs die out and do not grow into the twig or branch on which the spur occurs but occasionally the germs may continue into the bark of the branch at the base of the fruit spur and form a typical canker. Fruit spurs on the larger branches are a fruitful source of body infection and many cases of blight canker originate in this way.

Blight cankers may also originate through infection taking place at the tip of young actively growing branches, particularly if these be water sprouts on the larger branches, or trunks or through infection gaining a foothold in wounds of any kind. Such cankers on the larger limbs and trunk may retain active germs over the dormant season and form the so-called "hold-over" cankers which doubtless afford the only source of infection for the blossoms in the spring.

Appearance of the Disease.
The name "Fire Blight" is given to this disease because of the characteristic appearance of pear foliage on twigs or branches which have been killed by the blight organisms. The leaves turn black as though scorched by fire and frequently remain on the tree during the following winter. It should be noted that this color of the foliage is characteristic of the pear when it has been killed during the growing season. If a grower not familiar with the pear blight desires to know how the "twig blight" looks let him girdle a twig in mid-summer and watch the results. The foliage of the fruit spurs, killed by the blight, has much the same appearance as that of the twigs. On the apple the foliage of twig and fruit spur blight turns brown and dry.

The cankers are also quite characteristic but are very variable in appearance, depending on the age of the branch attacked and upon the condition of the tree. The disease progresses most rapidly in the fleshy outer layer of the bark and at first produces a watery appearance in the affected area. Later the tissues of the bark are more or less broken down and the cankers become dark in color and slightly sunken and are filled with a gummy substance which in active canker exudes. This exudate is at first sticky and contains myriads of live germs. It is attractive to insects which visit such cankers in great abundance and become covered with the organisms. If after becoming infected in this way they visit the blossoms for nectar they inoculate the blossom with the germs which find an easy access to the tissues of the blossom through the nectaries.

Besides the blight cankers found on the limbs and trunks one frequently finds in some varieties of pears and Spitzenberg apples a larger canker at the surface on the ground extending up on the trunk for some distance and down the large roots. This condition is called "collar rot" and may result from a blight canker appearing on the trunk and running down into the collar, or may result from infection taking place through twig blight running down water sprouts which come off at or near the surface of the ground. Wounds made by borers or gnawing animals may furnish the point of infection.

A foot rot may be caused by the organisms gaining entrance to the bark of roots through infection in the tips of water sprouts which come off some distance from the tree. This condition is common in some varieties of pears.

A pear tree when badly cankered is easily recognized at a distance in the early autumn by the general reddish cast to the foliage. When the foliage of only a part of the tree is reddish it indicates that a large canker has wholly or partly girdled one of the large branches. If the foliage of the whole tree is equally affected we should look for a canker at the collar or roots. It should be mentioned,

however, that any disorder of the root system, whether due to pear blight or some fungus or physiological disease will have a similar effect on the foliage. Whetzel has pointed out that when the apple is badly cankered, either on the larger limbs or trunk the foliage is pale green or grey, more or less dwarfed and curled, a condition which has been called by growers "mouse ear".

The pear and apple tree are covered by a protective layer of tissue, the cuticle, through which blight organisms cannot enter, except the nectaries and stigmas in the blossoms and actively growing tips of branches. Therefore except in cases of blossom infection and some cases of twig blight the germs can enter only through wounds which expose the sappy portion of the bark. Wounds may be caused in various ways—punctures of insects, birds (esp. suckers), gnawing of animals, bark or otherwise wounding by careless farm hands in cultivation, pruning or picking. Germs may also enter through growth cracks.

Method of Spreading the Disease.
The Fire Blight germs are naturally disseminated chiefly by insects at blossoming time as described above. Insects may also carry the germs to wounds on the branches or trunk. It is probable that a portion of the infection of the twigs resulting in twig blight may be brought about by insect punctures. The feet of birds may become infected with blight germs and the disease thus carried long distances. Sap suckers may "tap" an active canker and carry the germs to uninfected trees. Typical cases of body blight have been traced by Waite to this sort of infection. Rain may wash the germs from the ooze of active cankers to wounds, insect punctures or growth cracks lower down on the branches or trunk of the same tree.

One of the most fruitful sources of infection has been by the pruning shears or saw. In pruning, if an active canker is cut into, the tools become infected and serve as inoculating instruments to spread the disease throughout the orchard.

Infection may spread very rapidly from one district to another along highways where seedling apples and pears are allowed to grow unmolessted. Seedling trees springing from fruit cores thrown from car windows make of the great overland railroads a natural and easy road over which Fire Blight may travel for long distances through unsettled regions. This is one of the possible ways that the most favored valleys, naturally isolated by mountain barriers, may become infected.

Conditions Favoring the Disease.
The disease is more prevalent in orchards which are rapidly growing and in a succulent condition due to naturally rich soil, abundant cultivation or application of nitrogenous fertilizers. Heavy winter pruning has a tendency to stimulate rapid and succulent growth and renders the tree more subject to the attacks of the blight. Irrigation should be reduced to the minimum in infested orchards.

Method of Control.
The only known method of controlling Fire Blight is to cut out all cases of hold-over cankers wherever they appear. Spraying with fungicides is of only supplementary value and the various "blight cures" are worse than useless and it is a waste of time and money to apply them.

Experience has shown that it is of little permanent value to attempt to cut out the fruit spur and twig blight as they appear. Unless these forms of the disease extend into the branches on which they occur and a canker is formed the disease usually becomes naturally limited and the germs gradually die. Due to drying out of the canker so that at the beginning of the dormant season very few such cases show live germs present.

The efforts of the grower should be directed to cutting out all cases of blight canker and body canker during the fall, winter and early spring when the cankers have become more or less limited in their growth and are not actively spreading.

Summer cutting intelligently applied is frequently of great value particularly where there is only a little blight. In the autumn before the leaves fall is a good time to do the cutting as all cases of twig blight are easily observed and the condition of the foliage as already described, offers a convenient method of determining the presence of cases of root rot or collar rot and obscure cases of blight canker on body or lower limbs. Every orchardist in an infested district should examine all pear and apple trees at the time suggested and cut out all diseased portions. The trees should be particularly examined for cases of the collar rot. It is in this form of the disease that causes many trees to be killed outright.

In infested regions it has been found practical by the most successful owners of commercial orchards to keep one or more men in the orchard during the summer and fall whose sole duty it is to locate and cut out all cases of blight canker as soon as they appear. This work of inspection and eradication can very profitably be carried on during the summer and early autumn but the best time to do

the work is in the late fall, winter and early spring for reasons already given. If an orchard is gone over carefully by trained inspectors two or three times and all cases of this so-called "hold-over" blight be cut there will be no source of infection in the spring and consequently no blossom blight.

A final and very thorough inspection of orchards should be made in the early spring before the blossoms open so as to detect and remove any cankers which may have been missed in the earlier inspections. This is one of the most important inspections that should be given.

The matter of eradication is complicated, however, by the fact that insects may carry the disease considerable distances, and even though one grower may eliminate hold-over blight from his own orchard he may lose a large part of his crop by blossom blight and have his orchard re-infested by insects carrying the blight germs from hold-over cankers in a less careful neighbor's orchard. Apple and pear trees in town and city lots are seldom cared for in an up-to-date manner and in infested fruit districts become a very serious menace to commercial orchards in the vicinity.

How to Cut Out the Disease.
Where cankers occur on small limbs or are so extensive as to nearly girdle the limbs on which they occur the whole limb or branch should be removed. In cutting off such a limb it is important that one get well below the area of infection. For safety, the cut should be made at least a foot below apparent infection since in actively growing cankers the discoloration of the bark is frequently so slight as to be easily overlooked.

Where cankers occur on the larger limbs and trunks, if not too extensive, the limb or tree may frequently be saved by carefully shaving off the diseased bark of the cankers to the capwood. Here again one should cut well beyond the point of infection, especially above and below the canker since the germs spread most rapidly up and down from the point of infection.

In the pear at the beginning of the dormant period the diseased bark or hold-over cankers has a characteristic reddish tinge and with a little experience one can tell how far infection has occurred. In actively growing cankers in spring and summer it is frequently very difficult to determine the extent of the infection vertically as there may be very little discoloration. In apples it is usually more difficult to determine the extent of infection than in pears.

In cutting into the cankers it is convenient to use a gouge such as is used by carpenters. This is especially valuable where cankers occur in crotches of limbs. It has the special advantage of making a clean cut that can be easily sterilized.

Sanitary Measures.
In cutting out the cankers it is necessary that the tools be kept moist with some good disinfectant. If this is not done when cutting out an active canker each cut will re-inoculate the germs into the bark at the edges of the canker and the labor may thus be useless.

Corrosive sublimate in a solution of one part to one thousand of water has been found to be the most satisfactory disinfectant. It is convenient to buy this from the druggist in tablet form. The solution is a violent poison and should be so marked and carefully guarded. It should be kept in a glass bottle as it will corrode metal containers. It is convenient to carry a sponge or cloth which is kept moist with the disinfectant and used to wipe off the tools after each cut. After the process of cutting out a canker is completed sufficient time should be allowed for the canker to dry out thoroughly when the surface should be carefully disinfected, and then painted over with white lead paint. The wound should be kept painted with a coat of paint till healed over by the formation of callus.

When cutting out twig blight or when pruning out badly diseased limbs the same precaution should be taken and the pruning tools sterilized after each cut. All pruned branches should be immediately taken from the orchard and burned. This is especially important if cutting is done in summer.

In infested districts sterilization of all tools after each cut during regular summer or winter pruning should be made a regular practice since it might happen that a hold-over canker be cut into and the tools become infected and the disease thus spread throughout an orchard.

Supplementary Methods of Control.
Spraying with lime-sulphur solution—winter strength—after the final spring inspection and just before the buds open has been suggested by Waite as an aid in preventing the spread of Fire Blight. An application at that time would cover over any cases of hold-over canker and serve to disinfect the sticky ooze as well as render it less attractive to insects.

Waite's method of pruning the pear in vase form makes the best type of tree in which to fight the pear blight. Fruit spurs should be kept off the larger branches and under no condition should water sprouts be allowed to form on the trunk or from the roots. These when occurring on branches or trunks should be cut off flush with the wood. A convenient tool for this purpose is the carpenter's gouge. Should water sprouts spring up from the roots they should be cut off below the ground.

It is a good practice in infested districts to remove all "weed" trees of the Pome family which occur along roadsides or in the vicinity of commercial orchards as such trees are veritable nurseries for the blight.

Practicability of Controlling Fire Blight.
Fire Blight has proven so widespread and destructive and has ruined so many pear and apple orchards in various sections of the country that many persons have the erroneous idea that it can not profitably be controlled. In recent years it has been proven conclusively that where all the pear and apple growers in an infested

district work together, the control of Fire Blight is not only practicable but ceases to be a serious problem. The difficulty comes in getting all the growers to inspect their orchards and cut out the disease faithfully each year. There are always disinterested persons, and in Oregon at least, there are still some general farmers scattered through the fruit districts who own family orchards. It is difficult to make such men realize their responsibility in regard to Fire Blight. The town and city lot orchardist is in the same class. Efficient county fruit inspection can do a great deal toward persuading, and if necessary forcing, such persons to either keep the disease under control in their orchards or cut down the infested trees. Public sentiment must be aroused before the county fruit inspector can do efficient work. The greatest good comes in educating the people concerned, first to a realization of the danger of Fire Blight and then to the methods by which the disease may be kept under control. This work has been under the immediate direction of F. J. O'Bara, assistant pathologist in the department of agriculture. Through a system of county fruit inspection coupled with efficient education in regard to Fire Blight, public sentiment has been aroused and the people work together in controlling the disease.

It should be emphasized however, that the Fire Blight is a state and not a local problem. The disease is spreading and in time will without doubt occur in all parts of the state. It depends on the people whether or not it will be held under control. Complete eradication is not to be expected.

Note—It is proposed to issue from time to time circular bulletins dealing with crop pest problems. This circular is the first of the series. The department of entomology and plant pathology will be glad to answer inquiries relating to insect pests or plant diseases at any time. Always include with your inquiries as full a description of the trouble as possible and send specimens for examination. Address: Department of Entomology and Plant Pathology, Oregon Agricultural College, Corvallis, Oregon.

ECHO WOMAN IS STUNG BY SCORPION

PROMPT TREATMENT PREVENTS SERIOUS RESULTS

Mrs. Pete Sheridan Stung by Repitile—Echo Rebekah Visit Hermiston—Many go to Stanfield to Celebrate.

(Special Correspondence.)
Echo, Ore., July 4.—Mrs. Pete Sheridan of lower Butter creek, had the misfortune of being stung by a scorpion last Friday, but prompt attention and treatment prevented any serious results. The scorpion is rarely seen but it has been known that there are a few to be found in a part of this section.

Mr. and Mrs. Joe Bailey, Mr. and Mrs. C. R. Bonney, Mesdames Scholl, Reeves, Malcolm and Miss Laura Bonney all members of the Henrietta Rebekah lodge of this place went down to Hermiston Friday evening last and assisted in the installation of the officers of the lodge of that place. They spent a very pleasant evening and returned home on the motor next morning.

Miss Maybell Lambert and Miss Agnes Donahue left on Saturday evening for La Grande where they will visit for a few days.

Mrs. Sarah Thompson of Prineville, is here visiting with Mrs. Frank Spinning.
W. A. Jones returned Sunday morning from Portland where he has been the past week on business.
The Misses Nona and Vina Houser are visiting in Pilot Rock, having gone up on Saturday.
Mrs. Sarah Thompson of Prineville Oregon, is here visiting with Mr. and Mrs. J. Frank Spinning.
Asa Thomson and family have re-

SYRUP OF FIGS AND ELIXIR OF SENNA

CLEANSSES THE SYSTEM EFFECTUALLY; DISPELS COLDS AND HEADACHES DUE TO CONSTIPATION. BEST FOR MEN, WOMEN AND CHILDREN—YOUNG AND OLD.

TO GET ITS BENEFICIAL EFFECTS—ALWAYS BUY THE GENUINE. MANUFACTURED BY THE CALIFORNIA FIG SYRUP CO.

SOLD BY ALL LEADING DRUGGISTS ONE SIZE ONLY. REGULAR PRICE 50¢ A BOTTLE.

turned from their trip to Mexico. They report some very warm weather when they left for home.

Mrs. Jos. Bailey left here Saturday evening for Pendleton where she met a party of friends and went on to Baker City by automobile.

Jerome George of Worcester, Mass., arrived here this morning on a visit to his cousin F. F. George. Mr. George was here two years ago and spent his vacation, and could not forget the west. So we see him again.

Mrs. Ben Smith of Hermiston, was the guest of Mrs. M. C. Norman yesterday.

Mrs. J. E. Taylor made a short business trip to Pendleton this morning.

State of Ohio, City of Toledo, Lucas County, ss.

Frank J. Cheney makes oath that he is senior partner of the firm of F. J. Cheney & Co., doing business in the City of Toledo, County and State aforesaid, and that said firm will pay the sum of ONE HUNDRED DOLLARS for each and every case of Catarrh that cannot be cured by the use of Hall's Catarrh Cure.

FRANK J. CHENEY, Notary Public.

Sworn to before me and subscribed in my presence, this 6th day of December, A. D. 1888.

A. W. GLEASON, Notary Public.
Hall's Catarrh Cure is taken internally and acts directly on the blood and mucous surfaces of the system. Send for testimonials free.

F. J. CHENEY & CO., Toledo, O.
Sold by all Druggists, 75c.
Take Hall's Family Pills for constipation.

ENLARGE WAREHOUSE ON THE SNAKE RIVER

Lewiston Idaho.—The Interior Warehouse has begun the work of remodeling the big warehouse on the Snake river waterfront which it recently purchased.

The first work will be to excavate the ground on the lower deck in order to make the size of the first deck the same as the Snake river avenue floor which is known as the upper deck. The present structure is 50 by 250 feet in size and so soon as the warehouse portion is completed the plans will be perfected for building an additional 50 feet for the elevator and building in which will be installed the mix, cleaning and feed mill plant.

A Frightful Wreck

of train, automobile or buggy may cause cuts, bruises, abrasions, sprains or wounds that demand Bucklen's Arnica Salve—earth's greatest healer. Quick relief and prompt cure results. For burns, boils, sores of all kinds, eczema, chapped hands and lips, sore eyes or corns, it's supreme. Surest pile cure. 25c at Tallman & Co.

Austrian Alpinist's Tragic Fate.

Vienna.—An Austrian Alpinist named Wolfgang, while climbing in the Tyrol, was carried down a steep mountain by an avalanche and hurled into a pine tree. His feet caught in the branches and he was unable to free himself. His dead body was found hanging there.

Men's oxfords cheap at A. Eklund's.

Why Have an Overheated Kitchen in Summer?

When the sultry days come and the coal range makes the kitchen almost unbearable and cooking a dreaded task, put out the range fire and try the newest method of cooking in hot weather—use a

New Perfection Oil Cook-stove

What a contrast! The kitchen no longer is stifling hot, the work is now done with comfort, and the housewife is not worn out with the heat.



She saves her strength, keeps her health and is better able to enjoy the summer.

The New Perfection does everything that any other stove can do—all the family cooking, baking, washing and ironing. No smoke, no dust, no odor. Heat is applied directly and not wasted. A turn, and the flame is out.

The New Perfection stove has a Cabinet Top with shelf for keeping plates and food hot, drop shelves for the coffee pot or saucepans, and nickelled towel racks.

It has long turquoise-blue enamel chimneys. The nickel finish, with the bright blue of the chimneys, makes the stove very attractive and invites cleanliness. Made with 1, 2 and 3 burners; the 2 and 3-burner stoves can be had with or without Cabinet.

Every dealer everywhere, if not at yours, write for descriptive Circular to the nearest agency of the

Standard Oil Company (Incorporated)


Hay's Hair Health

Never Fails to Restore Gray Hair to its Natural Color and Beauty.

No matter how long it has been gray or faded. Promotes a luxuriant growth of healthy hair. Stops its falling out, and positively removes Dandruff. Keeps hair soft and glossy.

Will not soil skin or linen. Is not a dye. \$1 and 50c. bottles, at druggists. By mail 10c. Send 2c. for free book "The Care of the Hair Skin." Philadelphia Spec. Co., Newark, N.J., U.S.A. and Toronto, Ont., Canada.

Hay's Hairina Soap unequalled for the complexion, toilet and bath, red, rough, chapped hands. Keeps skin fine and soft. 25c. druggists. REFUSE ALL SUBSTITUTES. PENDLETON DRUG COMPANY.



Underwood Standard Typewriters

The Standard of typewriter manufacture, typewriter selling and typewriter work has been elevated by the advent and development of the UNDERWOOD

"The Machine You Will Eventually Buy"

Underwood Typewriter Company

(INCORPORATED)

68 Sixth Street - - Portland, Oregon